

Maternity – Document Immediate Post-op C-Section

This document details the documentation that occurs immediately post C-section delivery, when the mother is recovered on the Maternity ward.

Transfer the Patient

1. Transfer Patient from the OR (Operating Room) to appropriate location via Conversation Launcher. If Patient is on the Labour and Delivery OR, transfer patient using the rocket ship icon on the Tracking shell (Bed Transfer).



Powerplans

- Discontinue the Pre- Operative phase of the OB Cesarean Section Pre- Operative (Multiphase) Powerplan if not already done so by the Provider. Do not discontinue the Lab phase.
- 3. Initiate ANES Postpartum Intrathecal/Epidural Opioid and OB Postpartum Cesarean Section (Multiphase), Post Operative Powerplans.



Documentation in IView

- 4. In interactive View and I&O (iView), click on the OB Recovery and Postpartum Band
- 5. Click on the Phase 1 Arrival Section and document as applicable
- 6. Document vital signs in the Vital Signs Section under the OB Recovery and Postpartum Band
- 7. Document on Sedation in Sedation Scales in the Pasero Opiod Induced Sedation Scale



- 8. Continue further documentation in the OB Recovery Postpartum Band and chart in applicable sections eg Pain Modalities, Postpartum, Obstetrical Bleeding etc
- 9. When a patient is ready for discharge from the Postoperative Recovery area, calculate her discharge readiness by documenting in the Modified Aldrete Score (in the Sedation Scales section) as well as in the Discharge Criteria Section.

The Modified Aldrete Score and the Discharge Criteria Score both have score calculators . Double click in the cell to calculate the scores. Refer to the reference texts (by clicking on the blue reference text) to view the scores for discharge readiness.

🗙 OB Triage			Last 24 Hours
🗙 Antepartum			
🗙 Antenatal Testing	Find Item V Critical	High Low Abnormal	Unauth Flag
🗙 Labour and Delivery	Result	Comments Flag Date	Performed By
🗙 Newborn Delivery Data			
🗙 OB Recovery and Postpartum		01-Mar-20	118
PAIN ASSESSMENT		15:48 PS	Т
Comfort Measures	✓ Sedation Scales		
Obstetrical Bleeding	Sedation Scale Used	Sedation S	cale Used 🗙 🗙
Incision/Wound/Skin/Pin Site	4 Modified Aldrete Score	Modified 4	Aldrete Score
Surgical Drains/Tubes		Basero On	ioid Induced Sedation Scale
PSYCHOSOCIAL	Respiratory	Pasero Op	noid induced sedation scale
Parental Involvement	Circulation	Richmond	Agitation Sedation Scale
Activities of Daily Living	Level of Consciousness	University	of Michigan Sedation Scale
Urinary Catheter	♦ Movement	0	
IV Drips	0.000	~	
Anesthesia, OB	V SpO2	♦	
Specialty Gas Management	Modified Aldrete Score	\diamond	
Phase I Arrival			
Pain Modalities			
Sedation Scales			
Discharge Criteria			
Warming /Cooling	11		

Aldrete, Modified, Score		Reference				
CarePlan information	Aldrete, Modified, Score					
CarePlan information Chart guide Nurse preparation Patient education Policy						
odified Aldrete scoring	system (NAPAN, 2014)					
Catagory	Department of Status					
Category	Description of Status	Aldrete score				
Respirations	Able to deep breathe and cough freely	2				
	Dyspnea or limited breathing	1				
	Apneic	0				
O ₂ Saturation	Able to maintain O ₂ saturation > 92% on room air	2				
	Requires supplemental O ₂ to maintain SpO ₂ > 90%	1				
	O ₂ saturation < 90% even with supplemental O ₂	0				
Circulation	BP +/- 20% pre-op value	2				
	BP +/- 20-50% pre-op value	1				
	BP +/- > 50% pre-op value	0				
Level of Consciousness	Awake and oriented	2				
	Wakens with stimulation	1				
	Not responding	0				
Movement	Moves 4 limbs on own	2				
	Moves 2 limbs on own	1				
	Moves 0 limbs on own	0				
Discharge Criteria from O	ne to One Monitoring:					
The score for Respirations	must be 2.					
The score for O2 saturation	n must be 1 or greater .					
-						



×		11	
🗙 OB Recovery and Postpartum			
VITAL SIGNS	~	5.0 m	20-Eab-2019
Postpartum		R 📈	20-Feb-2018
Breast/Bottle Feeding			T1:09 PS1
Breastfeeding Assessment		Sedation Scales	
PAIN ASSESSMENT		⊿ Discharge Criteria	
Comfort Measures		Nausea and Vomiting	Controlled
Obstetrical Bleeding		Bleeding	Dressing si
Incision/Wound/Skin/Pin Site	2	p-i-	Dressing sim
Surgical Drains/Tubes		Pain	Controlled
PSYCHOSOCIAL		📟 Discharge Criteria Score	6
Parental Involvement			· · · · · · · · · · · · · · · · · · ·
Activities of Daily Living			
Urinary Catheter			
IV Drips			
Anesthesia, OB			
Specialty Gas Management			
Phase I Arrival			
Pain Modalities			
Sedation Scales			
Discharge Criteria			
Warming/Cooling			

Discharge Criteria Score

CarePlan information	Chart guide ONurse preparation	Patient education
Parameter	Description of Status	Score
Nausea and Vomiting	Controlled nausea/vomiting	2
	Further treatment required	1
	Uncontrolled nausea/vomiting	0
Bleeding	Dressing site dry and clean	2
	Hematoma present but not growing	1
	Increasing area of bleeding	0
Pain	Controlled pain	2
	Further treatment required	1
	Uncontrolled pain	0



Document Handoff

1. Document Handoff in Shift Report/Handoff tool and Transfer/Transport Section if appropriate.

Transfer

1. Transfer Patient to the appropriate location/bed using Conversation Launcher.